

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	77.8%
	Follow-up at Six Months	Expected	58.8%
	Response at Six Months	Expected	27.5%
	Remission at Six Months	Expected	17.7%
	Follow-up at 12 Months	Expected	58.8%
	Response at 12 Months	Expected	25.5%
	Remission at 12 Months	Expected	15.7%
Adolescent Mental Health and/or Depression Screening*		Average	98.5%
Colorectal Cancer Screening		Above Expected	79.7%
Optimal Asthma Control - Adults		Expected	47.8%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Expected	54.1%
HbA1c Control		Expected	77.0%
BP Control		Expected	87.8%
Daily Aspirin Use		Expected	98.0%
Statin Use		Expected	91.3%
Tobacco-Free		Expected	85.7%
Optimal Vascular Care		Expected	58.4%
BP Control		Expected	83.2%
Daily Aspirin Use		Expected	88.0%
Statin Use		Expected	94.4%
Tobacco-Free		Expected	78.4%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).