

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	83.4%
	Follow-up at Six Months	Expected	45.5%
	Response at Six Months	Expected	12.1%
	Remission at Six Months	Expected	5.5%
	Follow-up at 12 Months	Expected	45.5%
	Response at 12 Months	Expected	12.1%
	Remission at 12 Months	Below Expected	5.5%
Adolescent Mental Health and/or Depression Screening*		Below	81.5%
Colorectal Cancer Screening		Below Expected	63.5%
Optimal Asthma Control - Adults		Below Expected	5.0%
Optimal Asthma Control - Children		Below Expected	11.1%
Optimal Diabetes Care		Below Expected	37.0%
HbA1c Control		Expected	70.4%
BP Control		Below Expected	66.4%
Daily Aspirin Use		Expected	98.8%
Statin Use		Below Expected	83.9%
Tobacco-Free		Expected	82.7%
Optimal Vascular Care		Below Expected	44.6%
BP Control		Below Expected	68.9%
Daily Aspirin Use		Expected	88.7%
Statin Use		Expected	91.4%
Tobacco-Free		Below Expected	75.2%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).