

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	91.6%
	Follow-up at Six Months	Expected	56.8%
	Response at Six Months	Expected	22.2%
	Remission at Six Months	Expected	14.8%
	Follow-up at 12 Months	Expected	50.6%
	Response at 12 Months	Expected	23.5%
	Remission at 12 Months	Expected	14.8%
Adolescent Mental Health and/or Depression Screening*		Above	98.9%
Colorectal Cancer Screening		Above Expected	81.6%
Optimal Asthma Control - Adults		Expected	59.8%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Above Expected	65.0%
HbA1c Control		Above Expected	85.8%
BP Control		Above Expected	89.5%
Daily Aspirin Use		Expected	99.2%
Statin Use		Above Expected	96.6%
Tobacco-Free		Expected	86.6%
Optimal Vascular Care		Expected	65.8%
BP Control		Expected	87.1%
Daily Aspirin Use		Expected	89.6%
Statin Use		Expected	95.5%
Tobacco-Free		Expected	81.7%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).