

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	90.0%
	Follow-up at Six Months	Expected	46.4%
	Response at Six Months	Expected	18.6%
	Remission at Six Months	Expected	6.2%
	Follow-up at 12 Months	Expected	52.6%
	Response at 12 Months	Expected	21.1%
	Remission at 12 Months	Expected	13.4%
Adolescent Mental Health and/or Depression Screening*		Above	99.6%
Colorectal Cancer Screening		Below Expected	66.8%
Optimal Asthma Control - Adults		Above Expected	77.6%
Optimal Asthma Control - Children		Expected	62.1%
Optimal Diabetes Care		Above Expected	53.5%
HbA1c Control		Expected	77.3%
BP Control		Expected	85.3%
Daily Aspirin Use		Expected	99.5%
Statin Use		Expected	88.0%
Tobacco-Free		Expected	89.2%
Optimal Vascular Care		Expected	61.6%
BP Control		Expected	87.0%
Daily Aspirin Use		Expected	88.4%
Statin Use		Expected	91.8%
Tobacco-Free		Above Expected	89.0%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).