

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	91.3%
	Follow-up at Six Months	Above Expected	73.3%
	Response at Six Months	Expected	21.9%
	Remission at Six Months	Expected	13.3%
	Follow-up at 12 Months	Expected	61.9%
	Response at 12 Months	Expected	18.1%
	Remission at 12 Months	Expected	9.5%
Adolescent Mental Health and/or Depression Screening*		Below	81.6%
Colorectal Cancer Screening		Below Expected	48.5%
Optimal Asthma Control - Adults		-	-
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Below Expected	15.8%
HbA1c Control		Expected	64.9%
BP Control		Below Expected	50.9%
Daily Aspirin Use		Expected	100.0%
Statin Use		Expected	79.0%
Tobacco-Free		Expected	79.0%
Optimal Vascular Care		Expected	32.6%
BP Control		Below Expected	55.8%
Daily Aspirin Use		Expected	90.7%
Statin Use		Expected	90.7%
Tobacco-Free		Expected	62.8%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).