

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	93.6%
	Follow-up at Six Months	Above Expected	66.3%
	Response at Six Months	Expected	25.0%
	Remission at Six Months	Expected	13.1%
	Follow-up at 12 Months	Above Expected	65.0%
	Response at 12 Months	Expected	22.5%
	Remission at 12 Months	Expected	9.4%
Adolescent Mental Health and/or Depression Screening*		Below	81.2%
Colorectal Cancer Screening		Below Expected	65.2%
Optimal Asthma Control - Adults		Above Expected	61.3%
Optimal Asthma Control - Children		Expected	62.9%
Optimal Diabetes Care		Below Expected	39.1%
HbA1c Control		Expected	73.5%
BP Control		Below Expected	74.9%
Daily Aspirin Use		Expected	98.3%
Statin Use		Below Expected	85.4%
Tobacco-Free		Below Expected	81.4%
Optimal Vascular Care		Expected	53.2%
BP Control		Expected	78.1%
Daily Aspirin Use		Above Expected	91.1%
Statin Use		Expected	92.6%
Tobacco-Free		Below Expected	77.8%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).