

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	83.0%
	Follow-up at Six Months	Expected	43.3%
	Response at Six Months	Expected	16.7%
	Remission at Six Months	Expected	14.2%
	Follow-up at 12 Months	Below Expected	30.0%
	Response at 12 Months	Expected	15.0%
	Remission at 12 Months	Expected	10.0%
Adolescent Mental Health and/or Depression Screening*		Below	75.8%
Colorectal Cancer Screening		Below Expected	20.1%
Optimal Asthma Control - Adults		Below Expected	11.1%
Optimal Asthma Control - Children		Expected	24.6%
Optimal Diabetes Care		Expected	35.0%
HbA1c Control		Expected	60.5%
BP Control		Expected	76.9%
Daily Aspirin Use		Expected	99.6%
Statin Use		Expected	87.2%
Tobacco-Free		Above Expected	82.6%
Optimal Vascular Care		Expected	38.8%
BP Control		Expected	68.7%
Daily Aspirin Use		Expected	85.1%
Statin Use		Expected	97.0%
Tobacco-Free		Expected	65.7%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).