

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adolescent Mental Health and/or Depression Screening*		Below	85.1%
Colorectal Cancer Screening		Expected	65.7%
Optimal Asthma Control - Adults		Below Expected	28.3%
Optimal Asthma Control - Children		Below Expected	18.9%
Optimal Diabetes Care		Below Expected	32.8%
HbA1c Control		Expected	66.3%
BP Control		Below Expected	67.0%
Daily Aspirin Use		Expected	99.5%
Statin Use		Expected	86.5%
Tobacco-Free		Expected	80.7%
Optimal Vascular Care		Expected	45.0%
BP Control		Below Expected	67.6%
Daily Aspirin Use		Above Expected	91.9%
Statin Use		Expected	91.6%
Tobacco-Free		Expected	75.7%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).