

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	29.6%
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adolescent Mental Health and/or Depression Screening*		Average	93.0%
Colorectal Cancer Screening		Expected	67.3%
Optimal Asthma Control - Adults		Below Expected	0.0%
Optimal Asthma Control - Children		Below Expected	0.0%
Optimal Diabetes Care		Expected	43.1%
HbA1c Control		Expected	67.9%
BP Control		Expected	81.4%
Daily Aspirin Use		Expected	98.9%
Statin Use		Expected	85.0%
Tobacco-Free		Expected	88.7%
Optimal Vascular Care		Expected	61.3%
BP Control		Expected	81.1%
Daily Aspirin Use		Expected	91.5%
Statin Use		Expected	86.8%
Tobacco-Free		Expected	88.7%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).