

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	86.1%
	Follow-up at Six Months	Expected	54.6%
	Response at Six Months	Expected	19.9%
	Remission at Six Months	Expected	8.4%
	Follow-up at 12 Months	Expected	52.3%
	Response at 12 Months	Expected	20.2%
	Remission at 12 Months	Expected	9.9%
Adolescent Mental Health and/or Depression Screening*		Above	99.6%
Colorectal Cancer Screening		Above Expected	74.9%
Optimal Asthma Control - Adults		Above Expected	71.5%
Optimal Asthma Control - Children		Expected	66.3%
Optimal Diabetes Care		Expected	48.1%
HbA1c Control		Expected	73.5%
BP Control		Expected	83.4%
Daily Aspirin Use		Expected	98.8%
Statin Use		Expected	89.9%
Tobacco-Free		Expected	86.5%
Optimal Vascular Care		Expected	58.2%
BP Control		Expected	85.4%
Daily Aspirin Use		Expected	83.3%
Statin Use		Expected	93.9%
Tobacco-Free		Expected	85.4%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).