

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adolescent Mental Health and/or Depression Screening*		Average	95.7%
Colorectal Cancer Screening		Expected	71.2%
Optimal Asthma Control - Adults		Expected	55.2%
Optimal Asthma Control - Children		Below Expected	36.4%
Optimal Diabetes Care		Expected	52.1%
HbA1c Control		Expected	75.9%
BP Control		Above Expected	87.8%
Daily Aspirin Use		Expected	99.3%
Statin Use		Expected	89.1%
Tobacco-Free		Expected	86.9%
Optimal Vascular Care		Expected	61.2%
BP Control		Expected	85.1%
Daily Aspirin Use		Expected	88.0%
Statin Use		Expected	91.3%
Tobacco-Free		Expected	82.5%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).