

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	100.0%
	Follow-up at Six Months	Expected	48.6%
	Response at Six Months	Expected	16.2%
	Remission at Six Months	Expected	8.5%
	Follow-up at 12 Months	Expected	47.9%
	Response at 12 Months	Expected	21.8%
	Remission at 12 Months	Expected	13.4%
Adolescent Mental Health and/or Depression Screening*		Average	92.0%
Colorectal Cancer Screening		Above Expected	77.9%
Optimal Asthma Control - Adults		Below Expected	36.1%
Optimal Asthma Control - Children		Below Expected	30.0%
Optimal Diabetes Care		Expected	46.2%
HbA1c Control		Expected	75.0%
BP Control		Above Expected	87.5%
Daily Aspirin Use		Expected	97.7%
Statin Use		Expected	86.3%
Tobacco-Free		Below Expected	78.0%
Optimal Vascular Care		Expected	53.8%
BP Control		Above Expected	88.5%
Daily Aspirin Use		Expected	81.6%
Statin Use		Expected	92.1%
Tobacco-Free		Below Expected	74.7%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

## Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).