St. Francis Regional Medical Center

,



Clinical Quality Measures

2023 measurement year

| Measure | | Rating | Rate |
|---------------------------------|--|--------|--------|
| Adolescent Depression | PHQ-9 Utilization* | - | - |
| | Follow-up at Six Months | - | - |
| | Response at Six Months | - | - |
| | Remission at Six Months | - | - |
| | Follow-up at 12 Months | - | - |
| | Response at 12 Months | - | - |
| | Remission at 12 Months | - | - |
| | PHQ-9 Utilization* | - | - |
| ion | Follow-up at Six Months | - | - |
| Adult Depression | Response at Six Months | - | - |
| Jepi | Remission at Six Months | - | - |
| 봄 | Follow-up at 12 Months | - | - |
| Adı | Response at 12 Months | - | - |
| | Remission at 12 Months | - | - |
| Adole | scent Mental Health and/or Depression Screening* | - | - |
| Color | ectal Cancer Screening | - | - |
| Optimal Asthma Control - Adults | | - | - |
| Optin | nal Asthma Control - Children | - | - |
| | nal Diabetes Care | - | - |
| HbA10 BP Co | c Control ntrol | - | - - |
| Daily Aspirin Use | | - | - |
| Statin Use Tobacco-Free | | - | - - |
| Optimal Vascular Care | | - | - |
| BP Control | | - | - |
| Daily Aspirin Use Statin Use | | - | - |
| Tobacco-Free | | - | - |

⁻ Medical group not assigned to measure, did not have results, or did not meet reporting threshold

^{*}Measure not risk adjusted

St. Francis Regional Medical Center

Healthcare Effectiveness Data and Information Set (HEDIS) Measures

2023 measurement year

| Measure | Rating | Rate |
|---|---------|--------|
| Avoidance of Antibiotic Treatment in Acute Bronchitis/Bronchiolitis | Average | 72.90% |
| Breast Cancer Screening | Below | 72.20% |
| Cervical Cancer Screening | Average | 72.80% |
| Childhood Immunization Status (Combo 10) | - | - |
| Chlamydia Screening in Women | Average | 53.40% |
| Controlling High Blood Pressure | - | - |
| Diabetes Eye Exam | Below | 53.00% |
| Immunizations for Adolescents (Combo 2) | - | - |
| Follow-up Care for Children Prescribed ADHD Medication | - | - |
| Osteoporosis Management in Women Who Had a Fracture | - | - |
| Use of Spirometry Testing in the Assessment and Diagnosis of COPD | - | - |

⁻ Medical group not assigned to measure, did not have results, or did not meet reporting threshold

Ratings

Above: Medical group's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide/medical group rate (for unadjusted/HEDIS measures).

Average: Medical group's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide/medical group rate (for unadjusted/HEDIS measures).

Below: Medical group's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide/medical group rate (for unadjusted/HEDIS measures).

Total Cost of Care

2023 measurement year

| Population | Monthly Average Cost | Rating | Higher/Lower than Average |
|------------|-------------------------|--------|------------------------------|
| Overall | - | - | - |
| Adults | - | - | - |
| Pediatric | - | - | - |