



Dermatology *Partners*, Inc.

Consent for Treatment with Hyaluronidase

Hyaluronic acid (HA) fillers are sterile gels consisting of non-animal stabilized hyaluronic acid for injection into the skin to correct facial lines, wrinkles and folds, for lip enhancement and for shaping facial contours.

Occasionally these fillers need to be dissolved when the aesthetic treatment has not produced the desired outcome or there is a possibility of vascular occlusion or impending necrosis (tissue death) which could lead to compromise of healthy tissue.

Hyaluronidase is an enzyme which breaks down hyaluronic acid fillers, and is used to treat those problems. It can also break down naturally occurring hyaluronic acid present in the body and loss of volume and skin laxity which may not provide a good aesthetic result. It can take 24-48 hours for results to be seen. The treatment may need to be repeated.

Side effects of hyaluronidase can include allergic reaction. Seek emergency medical help if you have any of these signs of an allergic reaction: hives; difficulty breathing; swelling of your face, lips, tongue, or throat. Less serious side effects may include pain, itching, redness, or swelling where the medication was injected.

Other common injection-related reactions can occur. These include redness, swelling, pain, itching, bruising and tenderness at the injection site. These reactions typically resolve spontaneously within a few days .

I acknowledge that I will have to remain at the clinic for thirty minutes after the procedure so that I can be observed by the practitioner, and may need to return to the clinic in 1-2 weeks after treatment to assess if further hyaluronidase is to be administered.

Patient Signature

Print Name

Date

Provider Signature

Date