



# Vermillion Co Soil and Water Conservation District

Fall Fish Sale 2025

## Order Form



**Payment MUST accompany order**

Submit payment and order form to:

Vermillion Co SWCD

POB 86

Newport, IN 47966

765-492.3705

**Make Checks Payable: VCSWCD**

Name \_\_\_\_\_

Address \_\_\_\_\_

City, State, Zip \_\_\_\_\_

Phone: \_\_\_\_\_

Email: \_\_\_\_\_

**FISH DELIVERY: Thursday, October 23, 2024**

**Vermillion Co SWCD Office, Courthouse Annex Bldg, Newport from 10AM – NOON.**

*Fish will come pre-bagged for easier handling. Pickup reminder notices will be sent via email.*

SPECIES	SIZE	PRICE	QUANTITY	SUBTOTAL
Channel Catfish	4-6 inch	.85 each \$80/100	_____	\$ _____
Largemouth Bass	2-4 inch	1.55 each \$150/100	_____	\$ _____
Bluegill	2-3 inch	.80 each \$75/100	_____	\$ _____
Hybrid Bluegill	2-3 inch	.85 each \$80/100	_____	\$ _____
Redear Sunfish	2-3 inch	.90 each \$85/100	_____	\$ _____
Jumbo Bullfrog Tadpoles		.90 each \$55/100	_____	\$ _____
Fathead Minnows	1-3 lbs	\$15.50/lb	_____ lbs	\$ _____
	4+ lbs	\$13.00/lb	_____ lbs	\$ _____
Floating Fish Food	50 lb Bag	\$40.00	_____	\$ _____
Triploid Grass Carp (\$40 delivery fee)		\$17 each	_____	\$ _____
Black Crappie	3-4 inch	1.40 each \$135/100	_____	\$ _____

**Sub-Total** \$ \_\_\_\_\_

**7% Sales Tax** \$ \_\_\_\_\_

**Grass Carp Shipping** \$ \_\_\_\_\_

**Total** \$ \_\_\_\_\_

**DEADLINE TO ORDER ITEMS: Thursday, October 9 by 3 PM**

## **LARGEMOUTH BASS**

Largemouth bass are the primary predator in midwest bodies of water. They will feed on live fish and spawn every spring.

## **CHANNEL CATFISH**

Channel catfish are classified as a scavenger. Their diet ranges from live fish to crayfish, to commercial fish food. They will rarely reproduce in small ponds, unless provided with the appropriate structures.

## **BLACK CRAPPIE**

Black Crappie are a known predator whose dietary needs are equivalent to those of a largemouth bass. Black crappie typically do not perform exceptionally well in ponds less than 5 acres due to competition for live forage. Black crappie will spawn once a year.

## **TRIPLOID GRASS CARP**

Triploid Grass Carp are introduced into a pond to control or eliminate vegetation growth. The diet of a grass carp consists of a variety of species of vegetation.

## **STANDARD RECOMMENDATIONS**

600 BLUEGILL  
200 LARGEMOUTH BASS  
100-200 CHANNEL CATFISH  
100-200 REDEAR SUNFISH  
100-200 HYBRID BLUEGILL  
12-18 TRIPLOID GRASS CARP  
20# FATHEAD MINNOWS

- PER SURFACE ACRE -

## **REDEAR SUNFISH**

Redear Sunfish thrive on a diet that consists mainly of snails, worms, and insects. The diet of a redear makes them a wonderful aid in controlling or eliminating parasites within a body of water. Redear Sunfish will reach a mature size larger than that of a bluegill and will spawn once a year.

## **BLUEGILL**

Once mature, bluegill spawn 3 times a year. The prolific reproduction of the bluegill creates a substantial source of forage for the predator population of a body of water. We recommend a ratio of 3 bluegill to every 1 predator fish in order to sufficiently support the nutritional needs of the predator population.

## **HYBRID BLUEGILL**

Hybrid bluegill are the product of a female sunfish and a male bluegill. Hybrid bluegill reach a mature size larger than a straight bluegill and are known for being a more aggressive pan fish. Hybrid bluegill rarely reproduce due to the fact that roughly 80% of the population are males.

## **FATHEAD MINNOWS**

Fathead minnows are introduced into a pond as a forage source for predators. They rarely reach a size larger than 3" and will reproduce every 3 weeks.