



6828 Springfield Ave., Suite #1 | Laredo, Texas 78041

TO BE COMPLETED BY STAFF

Vaccine Given: _____

Lot: _____ Initials: _____

Date Given: _____

Patient Questionnaire for Flu Vaccination

PATIENT INFORMATION

Patient Name (Please Print Clearly)

Date of Birth

Age

PRE-VACCINE QUESTIONS

Are there any questions related to the flu vaccine that need to be discussed before the shot is given?

Please remember your visit today is with the nurse only. If you need to see a provider, you will need to make an appointment.

Yes

No

I understand that I can find the Vaccine Information Sheet for the Influenza Vaccine and Vaccine Side Effects Handout on our website www.drhcantu.net

Any immediate family member positive with COVID-19 or have been exposed to COVID-19?

Yes

No

Has any immediate family traveled out of the country or to COVID-19 hot spots?

Yes

No

If 12 and over, has patient received the COVID-19 vaccine in the past 2 weeks?

Yes

No

Is your child currently ill or running a fever, or had fever and respiratory symptoms in the past 72 hours? Any COVID-19 symptoms?

Yes

No

Please describe the illness:

HISTORY - Please check the box below if any of these items apply to your child

ALLERGY TO EGGS OR CHICKEN	REACTION TO FLU VACCINE IN THE PAST
DIABETES	IMMUNE COMPROMISED
PATIENT IS PREGNANT	RECEIVED A LIVE VIRUS VACCINE SUCH AS MMR, VARICELLA, OR ROTATEQ IN THE PAST 30 DAYS
ON LONG TERM ASPIRIN THERAPY	
SICKLE CELL DISEASE	ACUTE NEUROLOGICAL CONDITIONS OR HISTORY SEIZURES
OTHER CHRONIC RESPIRATORY OR CARDIAC DISORDERS, RENAL DYSFUNCTION	

SECOND DOSE INFORMATION

You must schedule a second dose if your child is under 9 years and has not previously received 2 doses of any type of flu vaccine before. The interval between the two doses must be at least 4 weeks.

I have scheduled my second dose

This is my second dose

No second dose needed

ACKNOWLEDGEMENT

Parent Signature

Date

Information reviewed and discussed by

Possible side effects of the flu vaccine include pain and redness at the site of injection. Less commonly, children may develop a fever or aches and pains, which last one to two days.