

Allergy Action Plan

Student Name: _____ Birth Date: _____
School: _____ Grade: _____ Teacher: _____

Place Student
Photo Here

ALLERGIC TO THESE ALLERGENS: _____

- Has Asthma (increases risk for severe reaction)
- Severe Allergy previously/suspected—Immediately give epinephrine & call 911— Start with Steps 2 & 3
- Mild Allergy – Itching, rash, hives – Give antihistamine, call school nurse and parent. Start with Step 1

► **STEP 1: IDENTIFICATION OF SYMPTOMS*** ◀

* Send for immediate adult assistance

Symptoms:

- If exposed to allergen, or allergen ingested, but **no symptoms**
- **Mouth** – Itching, tingling, or swelling of lips, tongue, mouth
- **Skin** – Hives, itchy rash, swelling of the face or extremities
- **Gut** – Nausea, abdominal cramps, vomiting, diarrhea
- **Throat** – Tightening of throat, hoarseness, hacking cough
- **Lung**** – Shortness of breath, repetitive coughing, wheezing
- **Heart**** – Faint, pale, blueness around mouth or nail beds, weak pulse, low B/P ..
- **Other**** – _____
- If reaction is progressing (several of the above areas affected) give

** Potentially life-threatening. – Note: The severity of symptoms can quickly change.

Type of Medication to Give:

(Determined by physician authorizing treatment)

<input type="checkbox"/> Epinephrine	<input type="checkbox"/> Antihistamine
<input type="checkbox"/> Epinephrine	<input type="checkbox"/> Antihistamine
<input type="checkbox"/> Epinephrine	<input type="checkbox"/> Antihistamine
<input type="checkbox"/> Epinephrine	<input type="checkbox"/> Antihistamine
<input type="checkbox"/> Epinephrine	<input type="checkbox"/> Antihistamine
<input type="checkbox"/> Epinephrine: Call 911	

► **STEP 2: GIVE MEDICATIONS** ◀

Epinephrine: inject intramuscularly (check one) EpiPen® EpiPen Jr®

- If Epinephrine is given, paramedics must be called! PROCEED TO STEP 3 BELOW.


Antihistamine/other: give _____ (Medication name & amount) by _____ (route/method)

- Notify parents and school nurse • Observe for increasing severity of symptoms • Call 911 as needed

IMPORTANT: Do NOT depend on asthma inhalers and/or antihistamines to replace epinephrine in a severe reaction.

EpiPen Directions:

- a. Pull off the GRAY Safety Cap
- b. Place BLACK TIP near OUTER-UPPER THIGH
- c. Swing and jab firmly until hearing or feeling a click
- d. Hold EpiPen in place **10 SECONDS**, remove, massage area
- e. Dispose of in red sharps container or give to paramedics



- The EpiPen can be injected through clothing.
- The individual may feel his/her heart pounding.
- This is a normal reaction to the medication.

► **STEP 3: EMERGENCY CALLS** ◀

1. **CALL 911** – Seek emergency care. State that an allergic reaction has been treated, and additional epinephrine may be needed.
2. Call Parents or Emergency Contacts

Parent completes Parent and Emergency Contact Names and Information below:

Parents/Emergency Contact Names: _____ Relationship: _____

Phone Number(s): _____

a. _____ 1.) _____ 2.) _____ () _____ () _____

b. _____ 1.) _____ 2.) _____ () _____ () _____

Parent/Guardian Signature

(Required)

Date _____

Physician completes form through Step 2

Physician Name (Printed) _____ Phone Number: _____ () _____

Physician Signature

(Required)

Date: _____

This form must be renewed annually or with any change in medication.

The Prescription Medication Authorization Form must be completed in addition to this Allergy Action Plan