



BOB WHITE
QUALITY FEED



44% Beef Builder Pellet

A high-protein medicated mixer pellet formulated for growing cattle on pasture or in dry lot situations

Features & Benefits:

- Support efficient weight gain, improve feed conversion, and overall growth performance
- Includes Rumensin® to promote weight gain while helping to control and prevent coccidiosis
- Contains urea for efficient rumen function to better utilize energy
- ClariFly® (Seasonal): An insect growth regulator that helps control fly populations by interrupting the development of larvae in manure

Feeding Directions: Feed 1-1.7 lbs per head/day.

IF FED MIXING WITH GRAIN:		Feed not less than 3.4, nor more than 24.6 lb of the above mix per head per day. Animals must have continuous access to the grain mix, do not let them run out.
Whole Shelled Corn	1775	
44% Beef Builde	225	
Total	2000	

IF FED WITHOUT MIXING WITH CORN: Feed 1.5 lb per head per day with ad libitum whole shelled corn. Limit roughage to 15% of the total diet. Animals must have continuous access to the whole shelled corn, do not let them run out.

GUARANTEED ANALYSIS:

Crude Protein*	(Min) 44.0%
Crude Fat	(Min) 3.4%
Crude Fiber	(Max) 3.0%
Calcium	(Min) 6.45%
Calcium	(Max) 7.70%
Phosphorous	(Min) 0.85%
Salt	(Min) 2.20%
Salt	(Max) 2.70%
Potassium	(Min) 2.3%
Selenium	(Min) 5 ppm
Vitamin A	(Min) 24,000 I.U./lb

* Includes not more than 18.00 % equivalent NPN.

Also available with ClariFly®

CAUTION: Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feed is safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing. **WARNING:** A withdrawal time has not been established for pruruminating calves. Do not use in calves to be processed for veal.

GERBER & SONS
— INC. —
FAMILY-OWNED SINCE 1905

Baltic, Ohio 43804
GerberAndSons.com