



BOB WHITE
QUALITY FEED

Elite-Calf Gro 18 Pellet

A premium grower pellet formulated to support optimal growth, strong muscles, and skeletal development in calves from post-weaning to six months of age

KEY INGREDIENTS: Midds, Fine Ground Corn, Soy Hulls, Molasses, Alfalfa Meal, Added Vitamins and Minerals



Features & Benefits:

- Unique blend of essential fatty acids that supports immune function, gut health, and nutrient absorption
- Precision prebiotic/postbiotic blend that supports immunity, gut health, and stress resilience
- ClariFly® (Seasonal): Feed-through fly control that minimizes fly emergence and promotes a cleaner, healthier environment
- Highly bioavailable trace minerals to support immunity, hoof health, and overall performance

Feeding Directions: Feed 6 to 10 lbs. per head daily, along with free-choice hay. Always ensure access to fresh, clean water.

Also available with ClariFly®.

Program Compatibility: Pairs well with Gerber and Sons Milk Replacer and Elite-Calf Starter 22 Tex.

GERBER & SONS
— INC. —
FAMILY-OWNED SINCE 1905

Baltic, Ohio 43804 / GerberAndSons.com

GUARANTEED ANALYSIS:

Medicated: For the prevention and control of coccidiosis due to *Eimeria zuernii* in calves (excluding veal calves).

Active Drug Ingredient:

Monensin (as monensin sodium) 38.7 g/ton

Crude Protein	(Min) 18.0%
Crude Fat	(Min) 4.8%
Crude Fiber	(Max) 11.5%
Acid Det. Fiber.	(Max) 13.0%
Calcium	(Min) 0.80%
Calcium	(Max) 1.30%
Phosphorous	(Min) 0.47%
Selenium	(Min) 0.4 ppm
Vitamin A	(Min) 10,000 I.U./lb.
Vitamin E	(Min) 36 I.U./lb.

CAUTION: Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feed is safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not feed undiluted. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not exceed the levels of Monensin recommended by the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

WARNING: A withdraw time has not been established for prurminating calves. Do not use in calves to be processed.