



BOB WHITE
QUALITY FEED

Elite-Calf Starter 22 Tex

A high-performance calf starter designed to give calves a strong foundation, enhancing immunity, growth, and development from day three through weaning for a smooth transition to solid feed

KEY INGREDIENTS: 31% Calf Pellet, Whole Corn, Whole Roasted Beans, Oats, Molasses, Cottonseed Hulls Meal



Features & Benefits:

- Unique blend of essential fatty acids that supports immune function, gut health, and nutrient absorption
- Precision prebiotic/postbiotic blend that supports immunity, gut health, and stress resilience
- ClariFly® (Seasonal): Feed-through fly control that minimizes fly emergence and promotes a cleaner, healthier environment
- Supports early rumen development and enhances feed intake for a robust daily gain

Feeding Directions: Feed free-choice to calves from day three through weaning. Ensure fresh, clean water is always available.

Also available with ClariFly®.

Program Compatibility: Pairs well with Gerber and Sons Milk Replacer and Elite-Calf Gro 18 Pellet.

GERBER & SONS
— INC. —
FAMILY-OWNED SINCE 1905

Baltic, Ohio 43804 / GerberAndSons.com

GUARANTEED ANALYSIS:

Medicated: For the prevention and control of coccidiosis due to *Eimeria zuernii* in calves (excluding veal calves).

Active Drug Ingredient:

Monensin (as monensin sodium) 44.3 g/ton

Crude Protein	(Min)	22.0%
Crude Fat	(Min)	5.1%
Crude Fiber	(Max)	7.5%
Acid Det. Fiber.	(Max)	9.5%
Calcium.	(Min)	0.50%
Calcium.	(Max)	1.00%
Phosphorous.	(Min)	0.46%
Selenium.	(Min)	0.4 ppm
Vitamin A	(Min)	11,000 I.U./lb.
Vitamin E	(Min)	57 I.U./lb.

CAUTION: Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feed is safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not feed undiluted. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not exceed the levels of Monensin recommended by the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

WARNING: A withdraw time has not been established for prurminating calves. Do not use in calves to be processed.