



BOB WHITE
QUALITY FEED

Opti-Calf Starter 20 Tex

A high-performance textured calf starter formulated to support early growth, immune function, and feed intake — laying the groundwork for long-term health and productivity

KEY INGREDIENTS: 31% Calf Pellet, Cracked Corn, Oats, Whole Roasted Beans, Molasses, Cottonseed Hulls Meal



Features & Benefits:

- Supports immune readiness, digestive health, and resilience to stress during early life transitions
- Rumensin® (Monensin): Enhances feed efficiency and helps prevent coccidiosis
- ClariFly® (Seasonal): An insect growth regulator that helps control fly populations by interrupting the development of larvae in manure
- Supports early feed intake, steady daily gain, and a smooth transition to dry feed

Feeding Directions: Offer free-choice feed to calves starting at three days old through post-weaning transition, and ensure clean, fresh water is always available.

Also available with ClariFly®.

Program Compatibility: Pairs well with Gerber and Sons Milk Replacers and Opti-Calf Gro 18 Tex.

GERBER & SONS
— INC. —
FAMILY-OWNED SINCE 1905

Baltic, Ohio 43804 / GerberAndSons.com

GUARANTEED ANALYSIS:

Medicated: For the prevention and control of coccidiosis due to *Eimeria zuernii* in calves (excluding veal calves).

Active Drug Ingredient:

Monensin (as monensin sodium) 45.0 g/ton

Crude Protein	(Min)	20.0%
Crude Fat	(Min)	4.6%
Crude Fiber	(Max)	8.0%
Acid Det. Fiber.	(Max)	9.5%
Calcium.	(Min)	0.40%
Calcium.	(Max)	0.90%
Phosphorous.	(Min)	0.43%
Selenium.	(Min)	0.2 ppm
Vitamin A	(Min)	6,000 I.U./lb.
Vitamin E	(Min)	19 I.U./lb.

CAUTION: Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feed is safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not feed undiluted. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not exceed the levels of Monensin recommended by the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

WARNING: A withdraw time has not been established for prurminating calves. Do not use in calves to be processed.