



BOB WHITE
QUALITY FEED

Select-Calf Gro 16 Tex

A high-quality textured grower feed formulated to support healthy development and steady weight gain in calves with a balanced energy-to-protein ratio and essential additives

KEY INGREDIENTS: 31% Calf Pellet, Cracked Corn, Oats, Molasses



Features & Benefits:

- A blend of whole grains and highly digestible fiber sources designed to promote growth and rumen development
- Supports a smooth transition from starter feed to full grower diets
- Rumensin® (Monensin): Enhances feed efficiency and helps prevent coccidiosis
- Promotes efficient gains and long-term productivity

Feeding Directions: Feed 6-10 lbs. per head per day along with free-choice hay and clean, fresh water.

Also available in an unmedicated blend.

Program Compatibility: Pairs well with Gerber & Sons milk replacers and Select-Calf Starter 18 Tex feed as part of a complete calf nutrition program.

GERBER & SONS
— INC. —
FAMILY-OWNED SINCE 1905

Baltic, Ohio 43804 / GerberAndSons.com

GUARANTEED ANALYSIS:

Medicated: For the prevention and control of coccidiosis due to *Elmeria zuernii* in calves (excluding veal calves).

Active Drug Ingredient:

Monensin (as monensin sodium) 39.0 g/ton

Crude Protein (Min)	16.0%
Crude Fat (Min)	3.0%
Crude Fiber (Max)	6.5%
Acid Det.. Fiber (Max)	7.5%
Calcium (Min)	0.35%
Calcium (Max)	0.85%
Phosphorous (Min)	0.41%
Selenium (Min)	0.2 ppm
Vitamin A (Min)	5,000 I.U./lb.
Vitamin E (Min)	16 I.U./lb.

CAUTION: Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feed is safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not feed undiluted. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not exceed the levels of Monensin recommended by the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

WARNING: A withdraw time has not been established for prurminating calves. Do not use in calves to be processed.