



BOB WHITE
QUALITY FEED



Step 2 Beef Mix

A balanced pellet-and-corn blend designed to take your cattle from 450lbs to market weight, with or without forage intake

KEY INGREDIENTS: Whole Corn, 44% Beef Builder Pellet, Molasses

Features & Benefits:

- An ideal ration to be used in steer stuffer and feedlot setting
- Delivers the right nutrition to support steady growth, efficient gain, and a healthy rumen function
- Formulated with whole shelled corn and our 44% Beef Boost Pellets to encourage intake
- Rumensin™ enhances feed efficiency and helps prevent coccidiosis
- Contains urea as an economical non-protein source

Feeding Directions: Begin feeding at 450lbs bodyweight and continue through to market weight. Provide 2.74-19.78 lbs per head daily, with access to clean, fresh water. Hay may be offered but is not required.

GERBER & SONS
— INC. —
FAMILY-OWNED SINCE 1905

Baltic, Ohio 43804 / GerberAndSons.com

GUARANTEED ANALYSIS:

Medicated: For increased rate of weight gain.

Active Drug Ingredient:

Monensin (as Monensin sodium) 36.4 g/ton

Crude Protein*	(Min)	12.0%
Crude Fat	(Min)	3.4%
Crude Fiber	(Max)	3.5%
Calcium	(Min)	0.65%
Calcium	(Max)	1.15%
Phosphorous	(Min)	0.6%
Magnesium	(Min)	0.2%
Copper	(Min)	23 ppm
Zinc	(Min)	80 ppm
Selenium	(Min)	0.7 ppm
Vitamin A	(Min)	5,000 I.U./lb.

*Includes not more than 2.55% equivalent NP.

CAUTION: Do not allow horses or other equine access to formulate containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle feed is safe for use in cattle and goats only. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions as reduced average daily gains may result. If feed refusal containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing. A withdrawal time has not been established for pre-ruminating calves. Do not use in. Calves to be processed for veal.

WARNING: Do Not feed to animals that are copper sensitive.