

Scoring the PHQ-9 modified for Teens

Scoring the PHQ-9 modified for teens is easy but involves thinking about several different aspects of depression.

To use the PHQ-9 as a diagnostic aid for Major Depressive Disorder:

- Questions 1 and/or 2 need to be endorsed as a "2" or "3"
- Need five or more positive symptoms (positive is defined by a "2" or "3" in questions 1-8 and by a "1", "2", or "3" in question 9).
- The functional impairment question (How difficult....) needs to be rated at least as "somewhat difficult."

To use the PHQ-9 to screen for all types of depression or other mental illness:

- All positive answers (positive is defined by a "2" or "3" in questions 1-8 and by a "1", "2", or "3" in question 9) should be followed up by interview.
- A total PHQ-9 score ≥ 10 (see below for instructions on how to obtain a total score) has a good sensitivity and specificity for MDD.

To use the PHQ-9 to aid in the diagnosis of dysthymia:

- The dysthymia question (In the past year...) should be endorsed as "yes."

To use the PHQ-9 to screen for suicide risk:

- All positive answers to question 9 as well as the two additional suicide items **MUST** be followed up by a clinical interview.

To use the PHQ-9 to obtain a total score and assess depressive severity:

- Add up the numbers endorsed for questions 1-9 and obtain a total score.
- See Table below:

Total Score	Depression Severity
0-4	No or Minimal depression
5-9	Mild depression
10-14	Moderate depression
15-19	Moderately severe depression
20-27	Severe depression

Next steps:

- If patient answers “No” to all questions 1 through 4, screening is complete (not necessary to ask question #5). No intervention is necessary (*Note: Clinical judgment can always override a negative screen).
- If patient answers **“Yes”** to any of questions 1 through 4, or refuses to answer, they are considered a **positive screen**. Ask question #5 to assess acuity:
 - ☐ **“Yes”** to question #5 = **acute positive screen** (imminent risk identified)
 - Patient requires a **STAT** safety/full mental health evaluation.
Patient cannot leave until evaluated for safety.
 - Keep patient in sight. Remove all dangerous objects from room. Alert physician or clinician responsible for patient’s care.
 - ☐ **“No”** to question #5 = **non-acute positive screen** (potential risk identified)
 - Patient requires a **brief** suicide safety assessment to determine if a **full** mental health evaluation is needed. **Patient cannot leave until evaluated for safety.**
 - Alert physician or clinician responsible for patient’s care.

Provide resources to all patients

- 24/7 National Suicide Prevention Lifeline 1-800-273-TALK (8255) En Español: 1-888-628-9454
- 24/7 Crisis Text Line: Text “HOME” to 741-741